

Torticollis and plagiocephaly in infancy: Therapeutic strategies

L. A. VAN VLIMMEREN¹, P. J. M. HELDERS², L. N. A. VAN ADRICHEM³, &
R. H. H. ENGELBERT²

¹Bernhoven Hospital, Veghel, The Netherlands, ²Wilhelmina Children's Hospital, Utrecht, The Netherlands,
³Erasmus University Medical Center Rotterdam, Sophia Children's Hospital, Rotterdam, The Netherlands

(Received 16 July 2004; accepted 20 September 2004)

Abstract

Background. Asymmetry in infancy is a diagnosis with a large spectrum of features, expressing an abnormal shape of parts of the body or unequal postures and movements, which might be structural and/or functional, with localized or generalized expression.

Purpose: The purpose of the present study is to highlight different therapeutic aspects of the most occurring asymmetries in infancy: congenital muscular torticollis, positional torticollis and plagiocephaly, based on best evidence in current literature.

Results: A flow chart is presented showing different pathways in therapeutic strategies, such as physical therapy, orthotic devices (helmet treatment and Dynamic Orthotic Cranioplasty) and surgery.

Conclusion: It is concluded that there are different views towards management on torticollis and plagiocephaly. A systematic therapeutic management to evaluate these asymmetries is indicated. The presented therapeutic flow chart might serve as a basis in order to achieve uniformity in therapeutic thinking and performance.

Keywords: *Asymmetry in infancy, conservative treatment, helmet therapy, physical therapy, plagiocephaly, torticollis*

Abbreviations: *CMT: Congenital muscular torticollis, DP: Deformational plagiocephaly, PT: Positional torticollis, SCM: Sternocleidomastoïd muscle*

La asimetría en la infancia es diagnóstica con un amplio espectro de características, expresando una forma anormal de partes del cuerpo o posturas y movimientos desiguales, las cuales podrían ser estructurales y/o funcionales, con una expresión localizada o generalizada. El propósito del presente estudio es remarcar los diferentes aspectos terapéuticos de las asimetrías más frecuentes en la infancia: torticollis muscular congénita, torticollis de posición y plagiocefalia, basado en la mejor evidencia de la literatura actual. Presentamos un flujo grama mostrando las diferentes vías de estrategia terapéutica, tales como la terapia física, los aparatos órtesicos (tratamiento con casco y Craneoplastia Ortésica dinámica) y cirugía.

Conclusión: Concluimos que hay distintos puntos de vista en cuanto al manejo de la torticollis y la plagiocefalia. Esta indicado un manejo terapéutico sistemático para evaluar estas asimetrías. El flujo grama terapéutico presentado podría servir como base para lograr uniformidad en el pensamiento y la aplicación de la terapia.

Introduction

Asymmetry in infancy is a descriptive diagnosis with a large spectrum of features, i.e. structural and/or functional, generalized or localized, regarding abnormal shape of parts of the body or unequal postures and movements, with a multi-factorial aetiological expression [1–7]. The appearance of asymmetry in spontaneous posture and movements

of the infant and an increase of the incidence of plagiocephaly without synostosis [2,4,5,8–11] is associated with the changed guidelines to prevent Sudden Infant Death (SID) [1,6,11–20].

The purpose of the present study is to highlight the different therapeutic regimes regarding congenital muscular torticollis, positional torticollis and plagiocephaly. The search strategy was focused on current peer-reviewed literature in *Medline*,

PubMed, *CINAHL* and *Cochrane*, with the keywords: asymmetry, infancy, torticollis, plagiocephaly, intervention, therapy and treatment. Related publications were also searched for in the references of all publications. No randomized controlled trials or systematic reviews were found. Non-controlled studies have different views concerning the treatment of asymmetry in infancy [20,21]. A review article concerning the diagnostic strategies for the evaluation of asymmetry in infancy has recently been published [6].

Asymmetrical features

Most occurring are generalized asymmetry in posture and movements [2,22–27] and localized asymmetries as congenital muscular torticollis, positional torticollis and plagiocephaly [6,22,28]. These disorders are causally heterogeneous symptoms of similar nosologic entities [6].

Torticollis, defined as localized asymmetry in infancy, with preferential posture of the head and asymmetric cervical movements, might be present at birth [22,29] or may develop in the first months of life as a result of an imbalance in the muscular function in the cervical region [30]. Secondary abnormalities of skull and muscles in the cervical region are associated.

Congenital muscular torticollis (CMT) is the type of torticollis with a unilateral contracture of the sternocleidomastoid muscle (SCM), often based on a pseudotumour of infancy [30–34]. Positional torticollis (PT) will develop in case of a persistent positional preference of the head, without evidence of morphologic changes in the SCM and may be induced by a deformational plagiocephaly at birth or/and a one-sided positioning after birth, during the first 1–5 months of life [29].

Deformational plagiocephaly (DP) [31,35,36] has been attributed to the sleeping position, congenital muscular torticollis or positional torticollis, neurological or cervical defects and premature birth [11,30,37–39] (Figure 1). The asymmetry of the head may be initiated pre-natally [29] and be exacerbated post-natally, when the child is laid in a supine position [21,40]. This type of plagiocephaly often firmly increases during the first weeks of life [30,36,41,42]. DP should be differentiated from craniosynostosis, which is the result of asymmetric premature closure of cranial sutures [43,44], apparently caused by inborn errors [45].

Localized defects may combine more or less generalized clusters of manifestations. Several synonyms referring to a generalized functional asymmetry point to abnormal position and shape of the head and face, scoliosis, rib cage molding, pelvic obliquity, as well as hip and foot asymmetry. The



Figure 1. Deformational plagiocephaly (Erasmus MC Rotterdam).

appearance of morphologic asymmetries is only a matter of time; any longstanding functional asymmetry will eventually result in a deformity [6].

Conservative strategies

Conservative strategies to intervene in positional torticollis, congenital muscular torticollis and deformational plagiocephaly are primarily physical therapy and helmet treatment [5,42,46,47] or Dynamic Orthotic Cranioplasty (DOC) [30,36,38,48–52]. No randomized clinical trials could be found in the literature. In general, however, conservative treatment seems to be beneficial when applied between ~2–8 months old infants [53].

Preventive counselling of parents on positioning, handling and nursing of the infant is important to minimize the risk of positional preference and to correct DP [1,10,19,54,55]. The content of the guidelines does not contradict with the recommendations on SID [1,54].

Whereas neonatal occipital flattening of the skull is a precursor to DP, Peitsch et al. [29] suggested an adjustment of the AAP recommendations to let children sleep in alternating head positions and sleeping in side-laying position. In order to stimulate the quantitative and qualitative motor development, it is recommended to place infants, when awake and under supervision, regularly in the prone position

('tummy time'), likely more than 5 minutes a day [4,8,10,16,19,54,56].

Physical therapy

No randomized controlled trials concerning physical therapy intervention and asymmetry in infancy could be found. However, there is agreement regarding the goals of treatment. Knowledge of the natural course of the asymmetry and differential diagnostics is essential when decisions about intervention have to be made [6].

Handling, positioning and movement therapy focus on active and passive symmetry in posture and movements. The first few months of life, a physiological asymmetry of the trunk is common, but has to disappear spontaneously before the first birthday [57]. Treatment of generalized asymmetry, without clear pathological signs or/and not combined with a localized problem, is not necessary before the age of 4 months, because of a physiological asymmetry possibly caused by neurological maturation [57]. Only handling and positional advices, to stimulate more symmetry in position and movements, are instructed to the parents [1,10,55]. Follow-up may be *mandatory*.

In the case of torticollis, range of motion in the cervical region has to be normalized, an eventually occurring SCM imbalance should be treated and the spontaneous positional and movement preferences should be minimized [19,30,58]. This will lead to symmetric motor performance and alignment, without structural impairments in range of motion or muscle function. The first 4–6 months of life intervention is expected to be most effective [21,30,41,53,58,59]. In case of the existence of a pseudotumour, a palpable mass centrally in the SCM related to CMT, physical therapy is indicated, even before 2 months of age, because of the negative influence on motor development. An increase of asymmetry may develop, due to fibrosis of the SCM, eventually resulting in structural asymmetry with deformational plagiocephaly.

Lying on the not flattened side of the occipital skull initiates natural remodelling of the skull [19]. The aim of physical therapy is to advise the parents about specific handling and positioning, but also to design a home treatment programme [19]. In recent publications, fair-to-excellent results of physical therapy interventions are reported; however, the studies were not randomized nor controlled and had small sample sizes [33,59–61].

There are several ways to increase range of motion. Passive stretching is mentioned, but the method is rarely explicated or explained [7,59]. Demirbilek and Atayurt [60] recommended passive and active

stretching of the SCM on the affected side in CMT, using firm pressure in both techniques. When therapy started before 3 months of age, the outcome was excellent. If therapy started within 3–6 months of age and within 6–18 months of age, 25% and 71%, respectively, had a fibrous contracture of the SCM muscle requiring myotomy. In a retrospective review of 277 patients with CMT, Binder et al. [41] described treatment strategy, divided in advices of positioning, handling and stretching under 3 months of age. The exercises focused on neck and trunk range of motion, equal weight bearing of the trunk and mid-line activities of the upper extremities, when older than 3 months of age. Several prospective, but non-controlled studies, by Cheng et al. [33,62,63] reported the good overall results of gentle manual stretching.

Taylor and Norton [59] advocated a programme to increase active range of motion and positioning to improve passive range of motion avoiding pain and resistance, with good-to-excellent outcomes in 96% of the children. The choice for this programme was based on the negative experiences with passive stretching.

Treatment should be focused on symmetric motor development and incremental active range of motion of the cervical spine. Passive manipulations or/and manual stretching in order to increase range of motion are obsolete, especially when pain is provoked, because it may cause micro-traumata in the soft tissues, eventually leading towards more fibrosis and consequent decrease in range of motion. Therefore, longstanding stretching with a low intensity without provoking pain is indicated to influence collagen structures and thereby range of motion [64,65].

Physical therapy contains extensive specific handling, positional advices and intensive correcting exercises regarding range of motion and movements towards symmetry, whereas passive manipulations, which provoke discomfort of the child, should be avoided.

Orthotic device; helmet treatment and dynamic orthotic cranioplasty

The natural history of the misshaped neonatal head is unknown, but in observing the heads of the adult population, it is obvious one could deduce a natural remodelling process [66]. DP may be treated with an orthotic device or the natural correcting growth may be expected [5,40,42,47,53].

The effort of an orthotic device is to use the remaining skull growth to redirect head shape, by allowing enough space in the helmet at the flattened areas. A molding helmet (Figure 2) is worn 15–22 hours per day and, after improvement



Figure 2. Remodeling helmet (Erasmus MC Rotterdam).

following 3–4 months of therapy, it is worn only at night [36]. Helmet treatment is generally recommended between 6–18 months of age [44]. Some authors mentioned the use of Dynamic Orthotic Cranioplasty (DOC) [30,36,38,48–52], by application of a dynamic band, which mildly pressures to the apexes of the frontal and occipital prominences, while creating voids over the adjacent areas so that growth of the normal areas is held constant. This treatment starts at 3–4 months of age [38]. The reason to indicate helmet treatment or DOC seems to be subjective, because the measuring methods are different and not always clearly described. No strict indications for this treatment are found. A uniform, easy applicable, valid and reliable measuring-instrument does not exist [6].

Two studies compared the influence of molding helmet and no-helmet periods on plagiocephaly [5,36]. However, the differences in rate of asymmetry of the skull between the two groups were very small [36]. Vles et al. [5] studied the effect of treatment of helmet vs. non-helmet in 105 infants with DP. The helmet treated group improved significantly better and faster, but were analysed only by a subjective cosmetic outcome score. Loveday and Chalain [42] compared orthotic helmets and active counterpositioning. Nevertheless, the intervention periods of both were very different, probably based on a lack of clear indicators. So,

conclusions are not possible. Other studies were not randomized nor controlled [38,52,66].

Surgery

Surgical treatment of a remaining less contractile SCM is generally indicated at the age of 12 months or later. If, in spite of physical therapy, there is a progressive decrease in contractility or range of motion of the SCM, differential diagnostics and surgical intervention may be considered at an earlier stage. Surgical procedures vary from simple open myotomy to radical resections of the SCM. Intensive post-operative physical therapy including scar treatment and procedures to remain full range of motion of the neck and to regain muscle length are routine for a period up to 4 months. At 2 years of age or older, surgical treatment is followed by an adjustable torticollis brace, to be worn for 3 months [32]. In rare cases, children may present with severe residual DP, which requires craniofacial surgery [19,67]. Craniosynostosis may be diagnosed by subsequently 3D-CT scanning and always requires craniofacial plastic surgery [45].

Flow chart therapeutic strategies

Based on the best available evidence in the current literature, different pathways in therapy of torticollis and plagiocephaly are presented in a flow chart. The algorithm (Figure 3) indicates the direction towards interventions and secondary differential diagnostics. The general view today is that postural and congenital muscular torticollis does need conservative intervention. The indication is related to age and range of motion.

In the first month, parents should be explained to, to prevent deformities and decrease in range of motion [19]. Evaluation of the asymmetry should be planned. If positional symmetry is reached at the age of ~3 months of age, intervention can be stopped. Asymmetry in position and movements with decreased range of motion indicates physical therapy [19]. If the child, at the age of 6 months, will not reach a full symmetric motor development and range of motion, differential diagnostics and continuing physical therapy is the best choice in intervention [33,59–61].

Persisting severe deformation of the skull at the age of 5–6 months requires specific attention; differential diagnostics which indicate orthotic device or follow-up. However, in the vast majority of cases a differential diagnosis is possible by means of clinical examination at an earlier follow-up (3–4 months of age). Radiological examination will help to identify the pathology. A delayed diagnosis could lead to a worsening of the prognosis. At 12 months of age,




Figure 3. Flow chart: therapeutic strategies in torticollis and plagiocephaly in infancy.

there is a final follow-up. When an obvious asymmetry in position and/or movements persists, possibly with any dysmorphism, diagnostics concerning possible vertebral column anomalies are indicated. Mainly cosmetic considerations will determine the outcome whether skull growth is acceptable or not.

Conclusions

Since no randomized clinical trials have been reported concerning therapeutic strategies and non-controlled studies have different views towards management on congenital muscular and positional torticollis and plagiocephaly, there is not only a great need for randomized controlled trials but also for a

structured approach of this problem. A flow chart was designed based on best available evidence in the literature regarding the therapeutic strategies in order to achieve uniformity in therapeutic thinking and performance.

Acknowledgements

The authors thank Marja AGC Schoenmakers, PCS, PT, PhD and Sonja Raaff of the Department of Pediatric Physical Therapy and Exercise Physiology, Wilhelmina Children’s Hospital/University Medical Center Utrecht, The Netherlands, for their contribution in the development of the diagnostic

flow chart and also Professor Yolanda van der Graaf, MD, PhD for her insightful comments and helpful suggestions in preparing this article.

References

- Boere-Boonekamp MM, van der Linder-Kuiper AT. Positional preference: prevalence in infants and follow-up after two years. *Pediatrics* 2001;107:339–343.
- Hamanishi C, Tanaka S. Turned head-adducted hip-truncal curvature syndrome. *Disability in Childhood* 1994; 70:515–519.
- Tunnessen W. Asymmetry. In: JS Freeman, S Robinson, Ml Fraser, editors. Signs and symptoms in pediatrics. Philadelphia, PA: JB Lippincott; 1988. p 509.
- Visscher F, van der Graaf T, Spaans M, van Lingen RA, Fetter WPF. Prone position favorable for motor development of infants. *Nederlands Tijdschrift voor Geneeskunde* 1998;142: 2201–2205.
- Vles J, van Zutphen S, Hasaart T, Dassen W, Lodder J. Supine and prone head orientation preference in term infants. *Brain Development* 1991;13:87–90.
- van Vlimmeren LA, Helders PJM, van Adrichem LNA, Engelbert RHH. Diagnostic strategies for the evaluation of asymmetry in infancy; a review. *European Journal of Pediatrics* 2004;163:185–191.
- Wei JL, Swartz KM, Weaver AL. Pseudotumor of infancy and congenital muscular torticollis: 170 cases. *Laryngoscope* 2001;111:688–695.
- Davis BE, Moon RY, Sachs HC, Ottolini MC. Effects of sleep position on infant motor development. *Pediatrics* 1998;102:1135–1140.
- Dewey C, Fleming P, Golding J, and the ALSPAC Study Team. Does the supine sleeping position have any adverse effects on the child? II. Development in the first 18 months. *Pediatrics* 1998;101:98.
- Hunt CE, Puczynski MS. Does supine sleeping cause asymmetric heads? *Pediatrics* 1996;97:127–129.
- Kane AA, Mitchell LE, Craven KP, Marsh JL. Observations on a recent increase in plagiocephaly without synostosis. *Pediatrics* 1996;97:877–885.
- American Academy of Pediatrics. Task force on positioning and SIDS. *Pediatrics* 1992;89:1120–1126.
- Dweyer T, Ponsonby A-L, Newman NM, Gibbons LE. Prospective cohort study of prone sleeping position and sudden infant death syndrome. *The Lancet* 1991;837: 1244–1247.
- Fleming PJ, Gilbert R, Azaz Y. Interaction between bedding and sleeping position in the sudden infant death syndrome; a population based case-control study. *British Medical Journal* 1990;301:85–89.
- de Jonge GA, Engelberts AC, Koomen-Liefting AJM, Kostense PJ. Cot death and prone sleeping position in the Netherlands. *British Medical Journal* 1989;298:722.
- Kattwinkel J, Brooks J, Keenan MJ, Malloy M. American academy of pediatrics; positioning and sudden infant death syndrome (SIDS): update. *Pediatrics* 1996;98: 1216–1218.
- Kattwinkel J, Brooks J, Keenan MJ, Malloy M. Changing concepts of sudden infant death syndrome: implications for the sleeping environment and sleep position. *Pediatrics* 2000;105:650–656.
- Mitchell EA, Engelberts AC. Sleeping position and cot deaths. *The Lancet* 1991;338:192.
- Persing JA. Prevention and management of positional skull deformities in infants. *Pediatrics* 2003;112:199–202.
- Rekate HL. Occipital plagiocephaly: a critical review of the literature. *Journal of Neurosurgery* 1998;89:24–30.
- Bridges SJ, Chambers TL, Pople IK. Plagiocephaly and head binding. *Archives of Diseases in Childhood* 2002;86: 144–145.
- Behrman RE, Kliegman RM, Jenson HB. Scoliosis, cranio-synostosis, torticollis. In: RE Behrman, RM Kliegman, HB Jenson, editors. Nelson textbook of pediatrics. 16th ed. Philadelphia, PA: W.B. Saunders Company; 2000. pp 1812–1813, 2082–2086, 2089–2091.
- Fulford GE, Brown JK. Position as a cause of deformity in children with cerebral palsy. *Developmental Medicine and Child Neurology* 1976;18:305–314.
- Lloyds-Roberts GC, Pilcher MF. Structural idiopathic scoliosis in infancy: a study of the natural history of 100 patients. *Journal of Bone and Joint Surgery* 1965;47B:520–523.
- Mau H, Gabe I. Die sogenannte Säuglingsskoliose und ihre krankengymnastische Behandlung. Stuttgart/New York: Georg Thieme Verlag; 1988.
- Palmén K. Prevention of congenital dislocation of the hip. The Swedish experience of neonatal treatment of hip joint instability. *Acta Orthopaedica Scandinavica Supplementum* 1984;55(suppl 208):58–67.
- Wynne-Davies R. Infantile idiopathic scoliosis: causative factors, particularly in the first six months of life. *Joint and Bone Joint Surgery* 1975;57:138–141.
- Boere-Boonekamp MM, van der Linden-Kuiper AT, van Es P. Preferential posture in infants; a serious call on health care. *Nederlands Tijdschrift voor Geneeskunde* 1997;141:769–772.
- Peitsch WK, Keefer CH, Labrie RA, Mulliken JB. Incidence of cranial asymmetry in healthy newborns. *Pediatrics* 2002;10:e72.
- Golden KA, Beals SP, Littlefield TR, Pomatto JK. Sternomastoid imbalance versus congenital muscular torticollis: their relationship to positional plagiocephaly. *Cleft Palate Craniofacial Journal* 1999;36:256–261.
- Bredenkamp JK, Hoover LA, Berke GS, Shaw A. Congenital muscular torticollis. *Archives of Otolaryngological Head and Neck Surgery* 1990;16:212–216.
- Cheng JCY, Tang SP. Outcome of surgical treatment of congenital muscular torticollis. *Clinical Orthopedics* 1999;362:190–200.
- Cheng JCY, Tang SP, Chen TMK. Sternocleidomastoid pseudotumor and congenital muscular torticollis in infants: a prospective study of 510 cases. *Journal of Pediatrics* 1999;134:712–716.
- Tang S, Liu Z, Quan X, Qin J, Zhang D. Sternocleidomastoid pseudotumor of infants and congenital muscular torticollis: fine-structure research. *Journal of Pediatric Orthopedics* 1998;18:214–218.
- Littlefield TR, Kelly KM, Pomatto JK, Beals SP. Multiple-birth infants at higher risk for development of deformational plagiocephaly: II. Is one twin at greater risk? *Pediatrics* 2002;109:19–25.
- Mulliken JB, van der Woude DL, Hansen M, LaBrie RA, Scott RM. Analysis of posterior plagiocephaly: deformational versus synostotic. *Plastic Reconstructive Surgery* 1999;103: 371–380.
- Argenta LC, David LR, Wilson JA, Bell WO. An increase in infant cranial deformity with supine sleeping position. *Journal of Craniofacial Surgery* 1996;7:5–11.
- Littlefield TR, Beals SP, Manwaring KH, Pomatto JK, Joganic EF, Golden KA, Ripley CE. Treatment of craniofacial asymmetry with dynamic orthotic cranioplasty. *Journal of Craniofacial Surgery* 1998;9:11–17.
- Turk AE, McCarthy JG, Thorne CH, Wisoff JH. The 'back to sleep campaign' and deformational plagiocephaly: is there

- cause for concern? *Journal of Craniofacial Surgery* 1996;7: 12–18.
40. Bruneteau RJ, Mulliken JB. Frontal plagiocephaly: synostotic, compensational or deformational. *Plastic Reconstructive Surgery* 1992;89:21–33.
 41. Binder H, Eng GD, Gaiser JF, Koch B. Congenital muscular torticollis; results of conservative management with long-term follow-up in 85 cases. *Archives of Physical Medicine and Rehabilitation* 1987;68: 222–225.
 42. Loveday BPT, de Chalain TB. Active counterpositioning or orthotic device to treat positional plagiocephaly? *Journal of Craniofacial Surgery* 2001;12:308–313.
 43. Fenichel GM. Disorders of cranial volume and shape. In: GM Fenichel, editor. *Clinical pediatric neurology*. Philadelphia, PA: WB Saunders Company; 1993. pp 375–378.
 44. Miller RI, Clarren SK. Long-term developmental outcomes in patients with deformational plagiocephaly. *Pediatrics* 2000;105:E26.
 45. Mathijssen IMJ. *Craniosynostosis: clinical and fundamental aspects*. Thesis Erasmus University Rotterdam. Rotterdam: Wyt Document Services; 2000.
 46. Clarren SK. Plagiocephaly and torticollis: etiology, natural history and helmet treatment. *Journal of Pediatrics* 1981;98:92–95.
 47. Pollack IF, Losken HW, Fasick P. Diagnosis and management of posterior plagiocephaly. *Pediatrics* 1997;99:180–185.
 48. Aliberti F, Pittore L, Ruggiero C, Cinalli G, Maggi G. The treatment of the positional plagiocephaly with a new thermoplastic orthotic device. *Child's Nervous System* 2002;18:337–339.
 49. Kelly KM, Littlefield TR, Pomatto JK, Manwaring KH, Beals SP. Cranial growth unrestricted during treatment of deformational plagiocephaly. *Pediatric Neurosurgery* 1999;30: 193–199.
 50. O'Broin ES, Allcut D, Earley MJ. Posterior plagiocephaly: proactive conservative management. *British Journal of Plastic Surgery* 1999;52:18–23.
 51. Ripley CE, Pomatto J, Beals SP, Joganic EF, Manwaring KH, Moss SD. Treatment of positional plagiocephaly with dynamic orthotic cranioplasty. *Journal of Craniofacial Surgery* 1994;5:150–159.
 52. Teichgraeber JF, Ault JK, Baumgartner J, Waller A, Messersmith M, Gateno J, Bravenec B, Xia J. Deformational posterior plagiocephaly: diagnosis and treatment. *Cleft Palate Craniofacial Journal* 2002;39:582–586.
 53. Carson BS, Munoz D, Gross G, Vanderkolk CA, James CS, Gates J, North M, McKnight M, Guarnieri M. An assistive device for the treatment of positional plagiocephaly. *Journal of Craniofacial Surgery* 2000;11:177–183.
 54. Hutchison BL, Thompson JMD, Mitchell EA. Determinants of nonsynostotic plagiocephaly: a case-control study. *Pediatrics* 2003;112:e316.
 55. Najarian SP. Infant cranial molding deformation and sleep position: implications for primary care. *Journal of Pediatric Health Care* 1999;13:173–177.
 56. Jantz JW, Blosser CD, Fruechting LA. A motor milestone change noted with a change in sleep position. *Archives of Pediatric and Adolescent Medicine* 1997;151: 565–568.
 57. Hadders M, Klip-Van den Nieuwendijk AWJ, Martijn A, van Eykern LA. Assessment of general movements: towards a better understanding of a sensitive method to evaluate brain function in young infants. *Developmental Medicine and Child Neurology* 1997;39:89–99.
 58. Engelbert R, Schoenmakers M, van Vlimmeren L. *Kinderorthopedie*. In: R Van Empelen, R Nijhuis-van der Sanden, A Hartman, editors. *Kinderfysiotherapie*. Maarssen: Elsevier Gezondheidszorg; 2000. pp 257–263.
 59. Taylor JL, Norton ES. Developmental muscular torticollis: outcomes in young children treated by physical therapy. *Pediatric Physical Therapy* 1997;9:173–178.
 60. Demirbilek S, Atayurt HF. Congenital muscular torticollis and sternomastoïd tumor: results of nonoperative treatment. *Journal of Pediatric Surgery* 1999;34:549–551.
 61. Ho BC, Lee EH, Singh K. Epidemiology, presentation and management of congenital muscular torticollis. *Singapore Medical Journal* 1999;40:675–679.
 62. Cheng JCY, Tang SP, Chen TMK, Wong MWN, Wong EMC. The clinical presentation and outcome of treatment of congenital muscular torticollis in infants—a study of 1,086 cases. *Journal of Pediatric Surgery* 2000;35: 1091–1096.
 63. Cheng JCY, Wong MW, Tang SP, Chen TM, Shum SL, Wong EM. Clinical determinants of the outcome of manual stretching in the treatment of congenital muscular torticollis in infants. A prospective study of eight hundred and twenty-one cases. *Journal of Bone and Joint Surgery* 2001;83-A: 679–687.
 64. Flowers KR, Lastayo P. Effect of total end range time on improving passive range of motion. *Journal of Hand Therapy* 1994;7:150–157.
 65. Glasgow C, Wilton J, Tooth L. Optimal daily total end range time for contracture: resolution in hand splinting. *Journal of Hand Therapy* 2003;16: 207–218.
 66. Moss SD. Nonsurgical, nonorthotic treatment of occipital plagiocephaly: what is the natural history of the misshapen neonatal head? *Journal of Neurosurgery* 1997;87: 667–670.
 67. Hansen M, Mulliken JB. Frontal plagiocephaly. *Journal of Craniofacial Surgery* 1994;21:543–553.

Copyright of Pediatric Rehabilitation is the property of Taylor & Francis Ltd. The copyright in an individual article may be maintained by the author in certain cases. Content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.